



(soh-VAHL-dee)

MOST IMPORTANT INFORMATION ABOUT SOVALDI

SOVALDI can cause serious side effects, including:

Hepatitis B virus reactivation: Before starting treatment with SOVALDI, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during or after treatment with SOVALDI. This may cause serious liver problems including liver failure and death. Your healthcare provider will monitor you if you are at risk for hepatitis B reactivation during and after you stop taking SOVALDI.

ABOUT SOVALDI

SOVALDI is a prescription medicine used with other antiviral medicines to treat chronic (lasting a long time) hepatitis C virus genotype 1, 2, 3, or 4 infection in adults. It is not known if SOVALDI is safe and effective in children under 18 years of age.

POSSIBLE SIDE EFFECTS OF SOVALDI

SOVALDI can cause serious side effects, including:

- **Those in the “Most Important Information about SOVALDI” section.**
- **Slow heart rate (bradycardia):** SOVALDI, when taken with amiodarone (Cordarone®, Nexterone®, Pacerone®), a medicine used to treat certain heart problems, may cause slow heart rate, which in some cases has led to death or the need for a pacemaker. Get medical help right away if you take amiodarone with SOVALDI and get any of the following symptoms:
 - fainting or near-fainting
 - extreme tiredness
 - dizziness or lightheadedness
 - shortness of breath
 - not feeling well
 - chest pains
 - weakness
 - confusion
 - memory problems

The most common side effects of SOVALDI when used in combination with ribavirin include:

- tiredness
- headache

And when used in combination with peginterferon alfa and ribavirin include:

- tiredness
- difficulty sleeping
- headache
- low red blood cell count
- nausea

These are not all the possible side effects of SOVALDI. Tell your healthcare provider if you have any new symptoms while taking SOVALDI.

IMPORTANT FACTS

This is only a brief summary of important information about SOVALDI and does not replace talking to your healthcare provider about your condition and your treatment.

BEFORE TAKING SOVALDI

Tell your healthcare provider about all of your medical conditions, including if you:

- have ever had hepatitis B virus infection
- have liver problems other than hepatitis C infection
- have severe kidney problems or you are on dialysis
- have HIV infection
- have any other medical condition
- are pregnant or breastfeeding, or plan to become pregnant or breastfeed
 - When taking SOVALDI with ribavirin or in combination with peginterferon alfa and ribavirin, you should also read those Medication Guides.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider.

SOVALDI and certain medicines may affect each other, or cause side effects.

GET MORE INFORMATION

- This is only a brief summary of important information about SOVALDI. Talk to your healthcare provider or pharmacist to learn more.
- Go to SOVALDI.com or call 1-844-READY41
- To learn about potential savings, call 1-855-7-MYPATH or go to SOVALDI.com

